



# Valproic acid induced encephalopathy – 19 new cases in Germany from 1994 to 2003 – A side effect associated to VPA-therapy not only in young children

Thorsten Gerstner<sup>a,\*</sup>, Deike Buesing<sup>a</sup>, Elke Longin<sup>a</sup>,  
Claudia Bendl<sup>a</sup>, Dieter Wenzel<sup>b</sup>, Brigitte Scheid<sup>b</sup>, Gisela Goetze<sup>b</sup>,  
Alfons Macke<sup>b</sup>, Gerhard Lippert<sup>b</sup>, Wolfgang Klostermann<sup>b</sup>,  
Geert Mayer<sup>b</sup>, Regine Augspach-Hofmann<sup>b</sup>, Sabine Fitzek<sup>b</sup>,  
Carl-Albrecht Haensch<sup>b</sup>, Markus Reuland<sup>b</sup>, Stephan A. Koenig<sup>a</sup>

<sup>a</sup> University Children's Hospital Mannheim, Neuropediatric Unit,  
Theodor-Kutzer-Ufer 1-3, 69167 Mannheim, Germany

<sup>b</sup> VPA-induced Side Effects Study Group: Children's Hospital Erlangen, Hamburg, Bruckmuehl,  
Hagen, Olpe, General Hospital Neukirchen, University Children's Hospital Jena, Helios-Klinikum  
Wuppertal, Neuwied, University Hospital Halle-Wittenberg, Germany

Received 23 January 2006; received in revised form 10 May 2006; accepted 23 May 2006

## KEYWORDS

Valproic acid;  
Side effects;  
Encephalopathy;  
Ammonia

**Summary** Valproic acid (VPA) is a broad-spectrum antiepileptic drug and is usually well-tolerated. Rare serious complications may occur in some patients, including haemorrhagic pancreatitis, bone marrow suppression, VPA-induced hepatotoxicity and VPA-induced encephalopathy. The typical signs of VPA-induced encephalopathy are impaired consciousness, sometimes marked EEG background slowing, increased seizure frequency, with or without hyperammonemia.

There is still no proof of causative effect of VPA in patients with encephalopathy, but only of an association with an assumed causal relation. We report 19 patients with VPA-associated encephalopathy in Germany from the years 1994 to 2003, none of whom had been published previously.

© 2006 British Epilepsy Association. Published by Elsevier Ltd. All rights reserved.

Sodium valproate (VPA) is a branched-chain fatty acid that is widely used in the therapy of epilepsy.

The anticonvulsive action of VPA has been demonstrated to be due to a combined pharmacological effect of increased  $\gamma$ -amino-butyric acid (GABA) levels<sup>1</sup> inhibition of *N*-methyl-D-aspartat (NMDA)

\* Corresponding author.

**Table 1** VPA-induced encephalopathies with hyperammonemia

	Age (year)	Sex	Underlying disease	Retar-ded	EEG	VPA-level (mg/l)	ASAT U/l, ALAT U/l,	Ammonia ( $\gamma$ mol/l) (11–55)	Add. AED	Add. disease	Outcome
1	65	M	Cerebri med. infarct	+	delta-focus	84.6	Normal	150	—	CHD, hypertonia	Recovery after withdrawal of VPA
2	4.6	F	Focal epilepsy	—	Generalized $\delta$ -activity	170	Normal	208	ETH	—	Recovery after withdrawal of VPA
3	1.9	F	Absence-epilepsy	—	Background slowing	<sup>a</sup>	38/33	111	—	Enteritis	Recovery after withdrawal of VPA
4	0.8	M	MELAS	+	Gen. $\delta$ -activity	<sup>a</sup>	4,—	258	PB, SUL, LTG,CL	Susp. of amino- acid metabolic disease	Recovery after withdrawal of VPA
5	2.6	M	Pseudo-Lennox- Gastaut	+	Background slowing	—	77, 36	159	TPM, ETH	Anaemia—	Recovery after withdrawal of VPA
6	16	F	IGE	—	Background slowing	74.5	Normal	261	—	—	Recovery after withdrawal of VPA

ALAT, alaninaminotransferase; ASAT, aspartat amino transferase; CHD, coronary heart disease; CL, clonazepam; ETH, ethosuximide; LTG, lamotrigine; MELAS, mitochondrial myopathy, encephalopathy, lactacidosis, stroke; OXC, oxcarbazepine; PB, phenobarbitone; SUL, sultiam; TPM, topiramate.

<sup>a</sup> Not done during encephalopathy.

**Table 2** VPA-induced encephalopathies without or slight hyperammonemia

	Age (year)	Sex	Underlying disease	Retarded	EEG	VPA-level (mg/l)	ASAT U/l, ALAT U/l,	Ammonia ( $\gamma$ mol/l) (11–55)	Add. AED	Outcome
1	52	M	Focal epilepsy	—	Theta-focus	63.6	—	Normal	CBZ, PB	Recovery after withdrawal of VPA
2	76	F	Cerebri med. infarct	+	Theta-focus	84.6	—	—	—	Recovery after withdrawal of VPA
3	77	F	Post-haemorrhage epilepsy	+	Background slowing	140	—	—	—	Recovery after VPA-reduction
4	28	M	Schizenzephalia	+	—	—	—	Normal	TPM, LEV, CL	Recovery after withdrawal of VPA
5	32	F	Epilepsy	—	Background slowing	75	—	Normal	CMZ	Recovery after withdrawal of VPA
6	41	M	Epilepsy	+	Background slowing, poly-spike-waves	—	Normal	Normal	PB, PHT	Recovery after withdrawal of VPA
7	63	F	Post-haemorrhage epilepsy	+	Background slowing	—	124, 28	—	PB, PHT	Recovery after withdrawal of VPA
8	64	M	Hypoxic brain damage	+	Background slowing	95.7	Normal	23	—	Recovery after withdrawal of VPA
9	74	F	Primary generalized epilepsy	—	Background slowing	95.2	23, —	—	—	Recovery after withdrawal of VPA
10	4.5	F	I GE	—	Background slowing	78	Normal	Normal	PB	Recovery after withdrawal of VPA
11	2.6	F	Bourneville Pringle	+	Background slowing	143.6	52, 436	68	OXC, LTG, CL	Recovery after withdrawal of VPA
12	72	F	Cerebri post. infarct	—	Delta-focus	—	Normal	64	—	Recovery after withdrawal of VPA
13	19	M	Focal epilepsy	—	Generalized $\delta$ -activity	72.1	Normal	69	—	Recovery after withdrawal of VPA

CBZ, carbamazepine; CHD, coronary heart disease; CL, clonazepam; IGE, idiopathic generalized epilepsy; PB, phenobarbitone; TPM, topiramate.

receptors<sup>2</sup> and a blockade of neuronal sodium channels.<sup>3</sup> VPA also effects a variety of different metabolic pathways and the compositing of membranes.<sup>4</sup>

Most of the side effects are mild and transient. However, more serious adverse reactions can occur such as hepatotoxicity,<sup>5</sup> encephalopathy,<sup>6</sup> coagulation disorders<sup>7</sup> pancreatitis<sup>8</sup> and bone marrow suppression.<sup>9</sup>

Three forms of encephalopathy in children and adults treated with VPA have been described: encephalopathy as direct toxic effect of VPA, with high serum levels of VPA but normal ammonia,<sup>10</sup> hyperammonemic encephalopathy<sup>11</sup> and encephalopathy with impaired liver function.<sup>12</sup> Encephalopathy was reported as an extremely rare complication of VPA therapy, described mostly in patients with inborn errors of metabolism, but also in patients without a known metabolic defect.<sup>13</sup>

Several publications showed an increased risk of encephalopathy when VPA was combined with topiramate.<sup>13,14</sup>


Direct drug effect on neurotransmitters is generally considered to be the main mechanism of VPA-induced encephalopathy.<sup>15</sup> VPA may lower glutamine synthesis via inhibition of glutamine synthase, thus contributing to hyperammonemia.<sup>16</sup> Another possible mechanism is direct neuronal toxicity induced by increased intracellular concentrations of glutamate and ammonium in astrocytes<sup>17</sup> which may lead to potential neuronal injury and perhaps cerebral edema.<sup>18</sup> It is not clear, whether changes in citrullinogenesis or in other steps of the urea cycle account for the increase of blood ammonia induced by valproate.<sup>19</sup> Up to 20% of asymptomatic patients treated with VPA may have mild hyperammonemia.<sup>20</sup>

We report 19 patients with VPA-induced encephalopathy in Germany from the years 1994 to 2003, none of whom had been published previously.

Since 1979, 51 cases of VPA-associated encephalopathy were reported in the literature.<sup>5,6,21–34</sup> Additionally, 13 cases of VPA-associated encephalopathy were reported in combination with TPM.<sup>7,13,14,35–37</sup>

## Methods

We sent a questionnaire asking for side effects of VPA therapy from 1994 to 2003, to all 1200 members (Paediatricians, Neurologists) of the "German Section of the International League against Epilepsy". We specially inquired about severe side effects such as hepatopathy, pancreatitis, encephalopathy and coagulation disorders. The physicians were also asked for the anonymized patient files in order



**Figure 1** Age distribution of VPA-induced encephalopathies.

to obtain details like VPA-dosage, laboratory parameters, the history and the outcome of the patients.

## Results

In the last 10 years altogether 19 cases of VPA-induced encephalopathy were reported in Germany by members of German Section of the International League against Epilepsy. We analyzed the medical files of these 19 patients. Another eight patients were reported but we were not able to study the complete files. These patients were therefore omitted.

Tables 1 and 2 present details of the 19 patients, divided into two subgroups: patients with encephalopathy and hyperammonemia and patients with normal serum-ammonia or slight hyperammonemia (69, 68 and 64  $\gamma\text{mol/l}$ ).

The average age of the patients was 38.6 years; the age distribution is shown in Fig. 1.


Eight of the patients were male, 11 female. Ten patients were mentally retarded, most of them children with infantile brain damage, only one had an inborn metabolic disorder (MELAS-syndrome). The VPA-treatment was started before the metabolic disorder was known.

All patients recovered after withdrawal of VPA. A 76-years-old women showed a prolonged onset with a slow recovery over 4 months. This patient was transferred to a nursery home when the encephalopathy began, but was able to live on her own again after the withdrawal of VPA. She additionally suffered from a Parkinson-syndrome, which may explain the prolonged normalisation.

VPA-induced encephalopathy may also occur at normal or slightly increased ammonia levels,<sup>38</sup> which was the case in 13 of our patients. Six patients, on the other hand, showed increased ammonia level. The average level was 191.2  $\gamma\text{mol/l}$  (111–259  $\gamma\text{mol/l}$ ) (Table 1).

## Discussion

VPA-induced encephalopathy is an important and severe side effect of AED-therapy. In contrast to



**Figure 2** Possible types of VPA-induced encephalopathy.

previous publications, not only young children with metabolic defects were affected in our series, but also adults.

Nevertheless, neurological disabilities and metabolic disorders were a risk factor for VPA-induced encephalopathy.

The following four types of VPA-induced encephalopathy have to be distinguished.

- (1a) Encephalopathy with normal ammonia in 13 patients in our study, a direct effect on neurotransmitters postulated.<sup>15</sup>
- (1b) Encephalopathy with hyperammonemia without liver failure in six patients in this study—an inhibition of urea cycle is considered to be the pathomechanism, but this could not be the only reason, because some patients show encephalopathy with a moderate hyperammonemia by inhibiting the urea cycle, some do not.
- (2a) Encephalopathy with hyperammonemia and liver failure. Within our questionnaire we have found nine cases of lethal hepatopathies, with increased ammonia levels (average 271  $\gamma$ mol/l) and signs of an encephalopathy (fatigue, nausea and EEG-slowness). Additionally, apathy, EEG-slowness and hyperammonemia was found in 5 of the 16 patients (31%) in the series of Scheffner and Koenig.<sup>39</sup>
- (2b) Encephalopathy without hyperammonemia but liver failure: 31 cases of reversible hepatotoxicity (with and without hyperammonemia). 4/31 cases showed normal ranges of ammonia but the above described signs of encephalopathy (not regarded here but in a separate paper) (Fig. 2).

For sure it is by all means possible to have the same pathophysiological mechanism leading to the encephalopathy 2a and 2b.

On the other hand, there are patients (up to 20–50% in the series of Laub, Altunbasak and Verrotti<sup>20,40,41</sup>) that have a slightly elevated serum ammonia without clinical abnormalities. This can be considered an effect of the inhibition of the urea cycle by VPA. Although, there is still a lack of evidence to this hypothesis. We do not routinely

measure ammonia in asymptomatic patients, as clinical consequences in these patients cannot be derived but patients and physicians are irritated without any reason.

From 1979 until now 51 patients with VPA-induced encephalopathy have been published in the literature. We found, however, 19 patients only in Germany in the last 10 years, in opposition to the statement, that VPA-associated encephalopathy is an extremely rare side effect. We therefore believe that the occurrence of this severe side effect is underestimated as reversible side effects are less likely to be published.

VPA-induced encephalopathy should also be considered in patients with rapidly progressing severe dementia who are treated with valproate. These patients should receive an EEG and blood tests for serum ammonia, liver enzymes and VPA-serum-level.

Fortunately all patients with VPA-induced encephalopathy but normal transaminases and normal coagulation can be expected to recover after withdrawal or reduction of VPA.

## References

1. Dodge JC, Illig AM, Snyder PJ, Badura LL. GABA levels within the medial preoptic area: effects of chronic administration of sodium valproic acid. *Psychoneuroendocrinology* 2000;**25**: 519–34.
2. Makarska-Bialek K, Kaminski RM, Czuczwar SJ. Influence of the antagonist of the glycine site of NMDA receptors, MRZ 2/576, on the anticonvulsant activity of conventional antiepileptic drugs in mice. *Pharmacol Rep* 2005;**57**:458–66.
3. Pugsley MK, Yu EJ, Mclean TH, Goldin AL. Blockade of neuronal sodium channels by the antiepileptic drugs phenytoin, carbamazepine and sodium valproate. *Proc West Pharmacol Soc* 1999;**42**:105–8.
4. Koenig St, Knolle J, Friedewald St, Koelfen W, Hannak G. The effects of valproic acid, carbamazepine and phenobarbitone on the fatty acid composition of erythrocyte membranes in children. *Epilepsia* 2003;**44**:708–11.
5. Koenig SA. Fatal liver failure associated with valproate therapy in a patient with Friedreich's disease: review of valproate hepatotoxicity in adults. *Epilepsia* 1999;**40**:1036–40.
6. Yehya N, Saldarini CT, Koski ME, Davanzo P. Valproate induced hyperammonemic encephalopathy. *J Am Acad Adolesc Psychiatry* 2004;**43**:926–7.

7. Teich M, Longin E, Dempfle CE, Koenig SA. Factor XIII deficiency associated with valproate treatment. *Epilepsia* 2004;**45**:187–9.
8. Batalden PB, Van Dyne BJ, Cloyd J. Pancreatitis associated with valproic acid therapy. *Pediatrics* 1979;**64**:520–2.
9. Acharya S, Bussel JB. Hematologic toxicity of sodium valproate. *J Pediatr Hematol Oncol* 2000;**22**:62–5.
10. Lheureux PE, Penalzoa A, Zahir S, Gris M. Science review: carnitine in the treatment of valproic acid-induced toxicity—what is the evidence? *Crit Care* 2005;**9**:431–40. [Epub 2005 Jun 10].
11. Cuturic M, Abramson RK. Acute hyperammonemic coma with chronic valproic acid therapy. *Ann Pharmacother* 2005;**39**:2119–22. [Epub 2005 Nov 15].
12. Rath A, Naryanan TJ, Chowdhary GV, Murthy JM. Valproate-induced hyperammonemic encephalopathy with normal liver function. *Neurol India* 2005;**53**:226–8.
13. Longin E, Teich M, Koelfen W, König SA. Topiramate enhances the risk of valproate associated side effects in three children. *Epilepsia* 2002;**43**:451–4.
14. Latour P, Biraben A, Polard E, Bentue-Ferrer D, Beauplet A, Tribut O, et al. Drug induced encephalopathy in six epileptic patients: topiramate? valproate? or both? *Hum Psychopharmacol* 2004;**19**:193–203.
15. Kwan P, Brodie MJ. Neuropsychological effects of epilepsy and antiepileptic drugs. *Lancet* 2001;**357**:216–22.
16. Duarte J, Macias S, Coria F, Fernandez E, Claveria LE. Valproate-induced coma: case report and literature review. *Ann Pharmacother* 1993;**27**:582–3.
17. Ricard C, Martin K, Tournier M, Begaud B, Verdoux H. A case of Parkinsonian syndrome, cognitive impairment and hyperammonemia induced by divalproate sodium prescribed for bipolar disorder. *Encephale* 2005;**31**:98–101.
18. Verrotti A, Trotta D, Morgese G, Chiarelli F. Valproate-induced hyperammonemic encephalopathy. *Metab Brain Dis* 2002;**17**:367–73.
19. Alonso E, Girbes J, Garcia-Espana A, Rubio V. Changes in urea cycle-related metabolites in the mouse after combined administration of valproic acid and an amino acid load. *Arch Biochem Biophys* 1989;**272**:267–73.
20. Laub MC. Hyperammonemia in valproate therapy in children and adolescents. *Nervenarzt* 1986;**57**:314–8.
21. Panda S, Radhakrishnan K. Two cases of valproate-induced hyperammonemic encephalopathy without hepatic failure. *J Assoc Phys India* 2004;**52**:746–8.
22. O'Neill M, Dubrey RW, Grocott-Mason RM. Valproate encephalopathy and hyperammonemia. *Postgrad Med J* 2002;**78**:316–7.
23. Verrotti A, Trotta D, Morgese G, Chiarelli F. Valproate-induced hyperammonemic encephalopathy. *Metab Brain Dis* 2002;**17**:367–73.
24. Osuna MT, Burcet J, Ramio L, Ustrell X, Silva Y, Molins A. Hyperammonemic encephalopathy associated with valproate acid. *Neurologia* 2003;**18**:347–50.
25. Elgudin L, Hall Y, Schubert D. Ammonia induced encephalopathy from valproic acid in a bipolar patient: case report. *Int J Psychiatry Med* 2003;**33**:91–6.
26. Lokrantz CM, Eriksson B, Rosen I, Asztely F. Hyperammonemic encephalopathy induced by a combination of valproate and pivmecillinam. *Acta Neurol Scand* 2004;**109**:297–301.
27. Payen C, Frantz P, Martin O, Parant F, Moulisma M, Pulce C, et al. Delayed toxicity following acute ingestion of valpromide. *Hum Exp Toxicol* 2004;**23**:145–8.
28. Yehya N, Saldarini CT, Koski ME, Davanzo P. Valproate-induced hyperammonemic encephalopathy. *J Am Acad Child Adolesc Psychiatry* 2004;**43**:926–7.
29. McCall M, Bourgeois JA. Valproic acid-induced hyperammonemia: a case report. *J Clin Psychopharmacol* 2004;**24**:521–6.
30. Reif A, Leonhard C, Mossner R, Lesch KP, Fallgatter AJ. Encephalopathy and myoclonus triggered by valproic acid. *Prog Neuropsychopharmacol Biol Psychiatry* 2004;**28**:1061–3.
31. Nakazato Y, Ando S, Yamamoto T, Tamura N, Shimazu K. Valproate-induced hyperammonemic encephalopathy in a patient with Sjogren's syndrome. *Rinsho Shinkeigaku* 2004;**44**:682–5.
32. Kimmel RJ, Irwin SA, Meyer JM. Valproic acid-associated hyperammonemic encephalopathy: a case report from the psychiatric setting. *Int Clin Psychopharmacol* 2005;**20**:57–8.
33. Hantson P, Grandin C, Duprez T, Nassogne MC, Guerit JM. Comparison of clinical, magnetic resonance and evoked potentials data in a case of valproic-acid-related hyperammonemic coma. *Eur Radiol* 2005;**15**:59–64.
34. Ricard C, Martin K, Tournier M, Begaud B, Verdoux H. A case of Parkinsonian syndrome, cognitive impairment and hyperammonemia induced by divalproate sodium prescribed for bipolar disorder. *Encephale* 2005;**31**:98–101.
35. Thygesen KS, Wolf P. Valproate-caused encephalopathy. *Ugeskr Laeger* 2005;**167**:3793–4.
36. Cheung E, Wong V, Fung CW. Topiramate-valproate-induced hyperammonemic encephalopathy syndrome: case report. *J Child Neurol* 2005;**20**:157–60.
37. Hamer HM, Knake S, Schomburg U, Rosenow F. Valproate-induced hyperammonemic encephalopathy in the presence of topiramate. *Neurology* 2000;**54**:230–2.
38. Solomon GE. Valproate-induced hyperammonemic encephalopathy in the presence of topiramate. *Neurology* 2000;**55**:606.
39. Koenig SA, Siemes H, Bläker F, Boenigk E, Groß-Selbeck G, Scheffner D, et al. Severe hepatotoxicity during valproate therapy: an update and report of eight new fatalities. *Epilepsia* 1994;**35**:1005–15.
40. Altunbasak S, Baytok V, Tasouji M, Herguner O, Burgut R, Kayrin L. Asymptomatic hyperammonemia in children treated with valproic acid. *J Child Neurol* 1997;**12**:461–3.
41. Verrotti A, Greco R, Morgese G, Chiarelli F. Carnitine deficiency and hyperammonemia in children receiving valproic acid with and without other anticonvulsant drugs. *Int J Clin Lab Res* 1999;**29**:36–40.